

Cal/Ecotox  
Exposure Factors for Bullfrog (*Rana catesbeiana*)\*

Page 1

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Age at Fledging, Metamorphosis, Weaning	1		6 - 7	mo	NR	NR	CA	a	1
Age at Fledging, Metamorphosis, Weaning			2 - 3	yr	B	Tadpole	NY	b	2
Age at Fledging, Metamorphosis, Weaning				yr	NR	Tadpole	KY	c	3
Age at Fledging, Metamorphosis, Weaning			3 - 6	mo	NR	Tadpole	AZ; CA	d	4
Age at Fledging, Metamorphosis, Weaning			13 - 16 or 24 - 25	mo	NR	Tadpole	MI	e	5
Age at Fledging, Metamorphosis, Weaning			24 - 27	mo	NR	Tadpole	IL	f	6
Age at Fledging, Metamorphosis, Weaning			12 - 14	mo	NR	Tadpole	KY	g	7
Age at Sexual Maturity	5		1 - 2	yr	F	Adult	MI	h	8
Age at Sexual Maturity				yr	F	Adult	CANADA	i	9
Age at Sexual Maturity				yr	M	Adult	MI	j	8
Age at Sexual Maturity				yr	M	Adult	CANADA	k	9
Age at Sexual Maturity			1 - 2	yr	NR	Adult	AZ; CA	l	4
Body Weight - Mean			0.46 - 0.57	lbs	NR		IL	m	6
Body Weight - Mean			0.01 - 0.10	lbs	NR		IL	n	6
Body Weight - Mean	0.46	8.16 SE	0.33 - 0.56	lbs	NR		IL	o	6
Body Weight - Mean	0.02			lbs	NR		IL	p	6
Body Weight - Mean	0.64		0.55 - 0.76	lbs	NR		IL	q	6
Body Weight - Mean	0.80		0.68 - 0.91	lbs	NR		IL	r	6
Body Weight - Mean	0.58		0.53 - 0.62	lbs	NR		IL	s	6
Body Weight - Mean	175.9			g	B	Adult	Lab	t	10
Body Weight - Mean			608 - 646	g	NR	Adult	Lab	u	11
Body Weight - Mean	142.8	77.4 SD	9.5 - 274.0	g	NR	Adult	CANADA	v	12
Body Weight - Mean	10.7		6.0 - 18.7	g	NR	Juvenile	Lab	w	13
Body Weight - Mean	29.8		18.5 - 51.6	g	NR	Juvenile	Lab	x	13
Body Weight - Mean	42.4		27.6 - 77.2	g	NR	Juvenile	Lab	y	13
Body Weight - Mean	17.5		13.1 - 41.6	g	NR	Juvenile	Lab	z	13
Body Weight - Mean	55.8		40.5 - 100.8	g	NR	Juvenile	Lab	aa	13
Body Weight - Mean			249 - 252	g	NR	NR	AR	ab	14
Body Weight - Mean		741.7 SE	15 - 200	g	NR	NR	CA	ac	1
Body Weight - Mean	see citation			g	NR	Tadpole	KY	ad	3
Body Weight - Mean			7 - 43	g	NR	Tadpole	MI	ae	5
Body Weight - Mean			0.5 - 5.4	g	NR	Tadpole	KY	af	7
Body Weight - Mean			20 - 36	g	NR	Tadpole	KY	ag	7
Clutch or Litter Size	7,360			#/egg mass	F	Adult	NJ	ah	15
Clutch or Litter Size			6,000 - 20,000	eggs/clutch	F	Adult	MI	ai	16
Clutches or Litters per year			1 - 2	clutches/yr	F	Adult	MI	aj	17
Clutches or Litters per year			1 - 2	clutches/yr	F	Adult	MI	ak	16
Dietary Composition	Review				B	Adult		al	18
Dietary Composition	Newly metamorphosed and larval Rana (87%), Aquatic snails (5%), Adult and larval aquatic Coleoptera (3%), Adult and larval Odonata (3%), Other (2%)				B	Adult	NM	am	19
Dietary Composition	crayfish (24%), wolf spiders (36%), long horned earwigs (20%), sowbugs (25%), german cockroaches (14%), crickets (16%), vascular plants (36%), inorganic material (12%), other animals (10%)				NR	Adult	AZ; CA	an	4

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Dietary Composition	see citation				NR	Adult	NY	ao	20
Dietary Composition	frogs (13.0%); shiners (27.5%); gastropods (16.1%); crayfish (16.3%); arachnida (4.5%); coleoptera adult (10.5%); hemiptera (13.5%); orthoptera (5.8%); lepidoptera larvae (4.5%); odonata adult (4.2%)			%	NR	Adult	MO	ap	21
Dietary Composition	frogs (12.1%); goldfish (30.3%); gastropods (12.4%); crayfish (9.3%); arachnida (4.4%); coleoptera adult (10.6%); hemiptera (8.2%); orthoptera (8.7%)			%	NR	Adult	MO	aq	21
Dietary Composition	Insecta (20.6%); Arachnida (4.8%); Amphibia (27.4%); Diploda (0.3%); Gastropoda (1.0%); Crustacea (4.3%);Teleosts (0.6%); Mammalia (5.0%); Oligochaeta (1.3%); Reptilia (6.4%); Pebbles and sand (8.0%); Vegetative material (12.3%); Digested material (7.3%)				B	Both Adult and Juv.	VA	ar	22
Dietary Composition	Insecta (51.8%); Arachnida (11.2%); Amphibia (2.8%); Diploda (1.3%); Gastropoda (1.3%); Crustacea (1.4%);Teleosts (1.1%); Mammalia (0.2%); Oligochaeta (1.3%); Reptilia (0.2%); Pebbles and sand (3.4%); Vegetative material (12.5%); Digested material (11.6%)				B	Both Adult and Juv.	VA	as	22
Dietary Composition	Coleoptera (2.3-6.5%), Wasps and bees (1.1-4.7%), Ants (0-3.5%), Homoptera (2.3-6.0%), Odonata (12.4-17.5%), Larval Lepidoptera (0.5-2.1%), Diptera (3.7-10.5%), Arachnids (0-3.2%), Miscellaneous terrestrial(10.5-11.1%), Hemiptera (19.2-23.2), Larval Odonata (3.4-5.0%), Larval Diptera (4.4-7.2%), Miscellaneous aquatic (1.9-7.4%), Frogs (5.1-14.5%), Metamorphosing Odonates (0-5.7%), Miscellaneous (1.8-7.3)				B	Both Adult and Juv.	MI	at	23
Dietary Composition	see citation				NR	NR	AR	au	14
Dietary Composition	Insecta (92.0%), Crustacea (13.7%), Arachnida (13.7%), Diploda (5.6%), Chilopoda (1.6%), Gastropoda (4.0%), Reptilia (3.2%), Amphibia (9.6%), Aves (1.6%), Osteichthyes (4.0%)				NR	NR	TX	av	24

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Dietary Composition	Coleoptera (43.6%), Notonectidae (10.3%), Diptera (6.6%), Hymenoptera (6.3%), Locustidae (6.0%), Ephemeroptera (4.3%), Gryllidae (4.3%), Protura (3.3%), Chironomidae (2.6%), Culicidae (2.0%), Lepidoptera (1.3%), Odonata (0.3%), Gerridae (0.3%), Mantidae (0.3%), Sepsidae (0.3%), Unidentified parts (21.3%), Decomposed tissue (18.0%), Spiders (16.0%), Rocks, grass, leaves (22.0%), Chitinous material (10%), Snails, Planorbid (9.0%), Frogs (5.6%), Snails, Physid (4.7%), Small fish (4.3%), Unknown vertebrates (2.3%), Tadpoles (1.3%), Salamander (1.0%), Feathers (1.0%), Worms (0.6%), Egg sack (0.3%), Rabbit pellet (0.3%), Peromyscus (0.3%)				NR	NR	CA	aw	1
Dietary Composition	Gastropoda (0.7%), Arachnida (0.2%), Crustacea (12.8%), Insecta (81.5%), Chilopoda (0.2%), Osteichthyes, Cyprinidae (8.7%), Ictaluridae (10.1%), Centrarchidae (2.9%), Amphibia (0.7%), Mammalia (0.5%)				NR	NR	OK	ax	25
Dietary Composition	Decapoda: Astacidae (47.7%); Lepidoptera (19.0%); Coleoptera: Lampryidae (5.8%); Chrysomelidae (5.8%); Carabidae (4.1%); Curculionidae (0.3%); Chilopoda (7.7%); sand, gravel (1.2%); unidentified (1%)			%	NR	NR	KY	ay	26
Food Ingestion Rate	3.3			%	NR	Juvenile	Lab	az	13
Food Ingestion Rate	5.9			%	NR	Juvenile	Lab	ba	13
Food Ingestion Rate	7.1			%	NR	Juvenile	Lab	bb	13
Food Ingestion Rate	4.0			%	NR	Juvenile	Lab	bc	13
Food Ingestion Rate	0.4			g	B	NR	VA	bd	22
Food Ingestion Rate	4.6			g	B	NR	VA	be	22
Food Ingestion Rate	2.33			g	B	NR	VA	bf	22
Growth Rate	0.1056	0.01427 SE		mm/d	B		CANADA	bg	9
Growth Rate	0.0653	0.01465 SE		mm/d	B		CANADA	bh	9
Growth Rate	0.1322	0.1109 SE		mm/d	F		CANADA	bi	9
Growth Rate	0.0930	0.02287 SE		mm/d	F		CANADA	bj	9
Growth Rate	0.1250	0.02155 SE		mm/d	F		CANADA	bk	9
Growth Rate	0.0466	0.02093 SE		mm/d	F		CANADA	bl	9
Growth Rate	0.0304	0.02256 SE		mm/d	F		CANADA	bm	9

# Exposure Factors for Bullfrog (*Rana catesbeiana*)

## Page 4

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Growth Rate	0.0449	0.00996 SE		mm/d	F		CANADA	bn	9
Growth Rate	0.1353	0.01079 SE		mm/d	M		CANADA	bo	9
Growth Rate	0.0841	0.01882 SE		mm/d	M		CANADA	bp	9
Growth Rate	0.0697	0.03720 SE		mm/d	M		CANADA	bq	9
Growth Rate	-0.0043	0.33534 SE		mm/d	M		CANADA	br	9
Growth Rate	0.0038	0.01654 SE		mm/d	M		CANADA	bs	9
Growth Rate			5 - 20	mm/yr	B	Adult	MI	bt	8
Growth Rate	review				B	Adult; Tadpole		bu	27
Growth Rate	0.4			stages/d	NR	Tadpole	Lab	bv	28
Growth Rate	0.02			stages/d	NR	Tadpole	Lab	bw	28
Home Range	8.6		2.0 - 37.1	ft	B	Adult	CANADA	bx	29
Home Range			> 80 - 90	ft	NR	Adult	MO	by	30
Inhalation Rate	see citation				B	Adult	Lab	bz	10
Longevity			8 - 10	yr	B	Adult	CANADA	ca	9
Longevity			7-8	yr	NR	Adult	CA	cb	31
Longevity			5 - 8	yr	NR	Adult	MI	cc	32
Metabolic Rate	see citation				B	Adult	Lab	cd	10
Metabolic Rate	53.3	2.1 SE		ul O2/g/hr	NR	Adult	Lab	ce	11
Metabolic Rate	15.5	1.2 SE		ul O2/g/hr	NR	Adult	Lab	cf	11
Metabolic Rate	1.1	0.1 SE		umol O2/g/hr	NR	Adult	Lab	cg	33
Metabolic Rate	24.56		18.3 - 36.6	ul O2/g/hr		NR	Lab	ch	34
Metabolic Rate	17.02		5.3 - 39.4	ul O2/g/hr		NR	Lab	ci	34
Metabolic Rate	29.36		22.5 - 33.2	ul O2/g/hr		NR	Lab	cj	34
Metabolic Rate	27.74		22.5 - 33.2	ul O2/g/hr	NR	NR	Lab	ck	34
Metabolic Rate	178.2	36.7		ul O2/g/hr	NR	Tadpole	Lab	cl	35
Metabolic Rate	5.9	0.3 SE		umol O2/g/hr	NR	Tadpole	Lab	cm	33
Metabolic Rate	6.9	0.5 SE		umol O2/g/hr	NR	Tadpole	Lab	cn	33
Metabolic Rate	see citation				NR	Tadpole	Lab	co	36
Metabolic Rate	see citation			ul/g/hr	NR	Tadpole	Lab	cp	37
Metabolic Rate	0.1582	0.009 SE		ml O2/hr/tadpole	NR	Tadpole	Lab	cq	38
Population Density			8.3 - 12.8	#/1000 ft2	B	Adult	CANADA	cr	29
Population Density	9.1			#/linear river km	NR	Adult	AZ; CA	cs	4
Population Density	90.9	97.7 SD	0 - 310	#/m2	B	Tadpole	Mendocino; CA	ct	39
Population Density	18.2	13.1 SD	1 - 52	#/m2	B	Tadpole	Mendocino; CA	cu	39
Population Density			0.9 - 13.2	#/m2	NR	Tadpole	KY	cv	3
Surface Area	Log SA = 0.923 + (0.711Log BW)			cm2	NR	Adult	Lab	cw	40
Surface Area	SA = 0.953W^0.725			cm2	NR	NR		cx	34
Surface Area	15.09	0.85 SE		cm2	NR	Tadpole	Lab	cy	33
Survival/ Mortality	0.42				M		MI	cz	41
Survival/ Mortality			8 - 21	%	B	Adult	MI	da	8
Survival/ Mortality	0.23				M	Adult	MI	db	41

Endpoint Type	Endpoint Value	Error	Range	Units	Sex	Life Stage	Location	Note	Reference
Survival/ Mortality	0.52				M	Adult	MI	dc	41
Survival/ Mortality			11.8-17.6	%	NR	Tadpole	KY	dd	3
Territory Size	17.8	5.9 SD		ft	M	Adult	MI	de	42
Territory Size	4.1			m	M	Adult	NJ	df	15
Time of Fledging or Metamorphosis	Review				B	Tadpole		dg	18
Time of Fledging or Metamorphosis	July - Oct.				B	Tadpole	NY	dh	2
Time of Fledging or Metamorphosis	July - Sept.				NR	Tadpole	KY	di	3
Time of Fledging or Metamorphosis	Aug. - Oct.				NR	Tadpole	AZ; CA	dj	4
Time of Fledging or Metamorphosis	Aug. - Oct. or Mar. - Apr. of following yr.				NR	Tadpole	AZ; CA	dk	4
Time of Fledging or Metamorphosis	July - Sept.				NR	Tadpole	MI	dl	5
Time of Fledging or Metamorphosis	June (begin), July - Aug. (peak), Oct. (end)				NR	Tadpole	KY	dm	7
Time of Fledging or Metamorphosis	May (begin), July-Aug. (peak), Oct. (end)				NR	Tadpole	MO	dn	30
Time of Mating/ Laying	Review				B	Adult		do	18
Time of Mating/ Laying	May - June				F	Adult	IL	dp	6
Time of Mating/ Laying	Apr - June				F	Adult	NJ	dq	15
Time of Mating/ Laying	May (begin), June (peak), Aug. (end)				F	Adult	MO	dr	30
Time of Mating/ Laying	July				F	Adult	Mendocino; CA	ds	39
Time of Mating/ Laying	early Apr. (begin), late Apr. - early May and late May - early June (peaks), July (end)				NR	Adult	AZ; CA	dt	4
Time of Torpor or Hibernation	Oct. - Mar.				NR	Adult	MO	du	30
Time of Torpor or Hibernation	Oct. - April				NR	Adult	OH	dv	43
Time of Torpor or Hibernation	Oct. - Mar.				NR	Tadpole	IL	dw	6
Water Ingestion Rate	0.18	0.02 SE		ml/100 g/hr	NR	Tadpole	Lab	dx	44

- Notes**
- a time to complete from egg through metamorphosis; N=300; San Joaquin Experimental Range, O'Neals
  - b N=NR; Ithaca
  - c N=NR; Coldstream Pond and Fred Pond, northeast of Lexington, [38°06'23"N, 84°30'08"W]
  - d time to metamorphosis from first breeding clutch; N=NR; lower Colorado River between Laguna and Morelos Dams
  - e Referred to as "age at transformation"; range of snout-vent length at transformation = 35-57 mm; N=1051; Crane Pond, E.S. George Reserve, Livingston County [lat., 42.5N]; See paper for review of larval life history characteristics as a function of latitude
  - f N=NR; Ridge Lake, Charleston, Coles County
  - g Time to transformation after hatching from eggs; N=3500; ponds, University of Kentucky, Lexington [lat., 38°06'23"N; long., 84°30'08"W]
  - h age after metamorphosis when produced eggs; N=NR; Crane Pond, E.S. George Reserve
  - i Average age at maturity, measured in years post-transformation; N=567; Algonquin Provincial Park, Ontario [lat., 45°35'N; long., 78°30'W]
  - j age after metamorphosis; development of secondary sex characteristics; N=NR; Crane Pond, E.S. George Reserve
  - k Average age at maturity, measured in years post-transformation; N=567; Algonquin Provincial Park, Ontario [lat., 45°35'N; long., 78°30'W]
  - l N=NR; lower Colorado River between Laguna and Morelos Dams
  - m N=5; Age=end of 3rd yr; Ridge Lake, Charleston, Coles County
  - n N=5-13; Age=during 1st yr; May - Aug.; Ridge Lake, Charleston, Coles County
  - o N=5; Age=end of 2nd yr; Ridge Lake, Charleston, Coles County
  - p at metamorphosis; N=48; Age=at metamorphosis; Ridge Lake, Charleston, Coles County
  - q N=6; Age=end of 5th yr; Ridge Lake, Charleston, Coles County

r	N=5; Age=end of 6th yr; Ridge Lake, Charleston, Coles County
s	N=3; Age=end of 4th yr; Ridge Lake, Charleston, Coles County
t	N=30
u	range of mean body weights; N=8
v	N=39; Apr. - Sept.; York County, New Brunswick [lat., 45°o47'; long., 66°o37']
w	N=25; Age=newly transformed
x	N=25; Age=2 mo post-transformation
y	N=25; Age=3 mo post-transformation
z	N=25; Age=1 mo post-transformation
aa	N=25; Age=4 mo post-transformation
ab	range of mean body weights; N=29 - 62; Mar. - June; streams, ponds, strip pits in Pulaski, Loanoke and Saline counties
ac	N=300; San Joaquin Experimental Range, O'Neals
ad	figure of annual variation in weight; N=27 - 1817; Coldstream Pond and Fred Pond, northeast of Lexington, [38o06'23"N, 84o30'08"W]
ae	See paper for weight to developmental stage relationship (Gosner stages 27-40); N=120; May; Crane Pond, E.S. George Reserve, Livingston County [lat., 42.5N]; See paper for annual variation in body size at metamorphosis
af	Range of means; N=41 - 194; Age=Taylor-Kollros stages II - IX; July - Oct.; ponds, University of Kentucky, Lexington [lat.,38°o06'23"N; long., 84°o30'08"W]; See citation for complete stage to weight relationship
ag	Range of means; N=23 - 235; Age=Taylor-Kollros stages XVIII - XXI; July - Aug.; ponds, University of Kentucky, Lexington [lat.,38°o06'23"N; long., 84°o30'08"W]; See citation for complete stage to weight relationship
ah	Number of eggs in a 10x10cm area used to estimate eggs per mass; N=36; April - June; Basking Pond, Great Swamp National Wildlife Refuge, Morris County
ai	N=NR; June - July; Crane Pond, E.S. George Reserve
aj	Double clutching observed in 5 out of 73 females; N=73; May - June
ak	N=NR; Crane Pond, E.S. George Reserve
al	N=NR; Reviews literature on foods of adult bullfrogs
am	Total percent volume of stomach contents, frog snout-vent length range = 90 - 190 mm; N=138; July; Wall Lake, southeast Catron County
an	% frequency of occurrence in stomach contents; N=100; July - Sept.; lower Colorado River between Laguna and Morelos Dams
ao	% frequency of 26 food items in stomach contents (family level); N=24; July; Cranberry Lake Biological Station, Barber Point, Cranberry Lake
ap	% of stomachs containing major (>4%) food items; N=553; yr-round; bait minnow pond, Richland; see citation for detailed list of prey items and seasonal changes in diet
aq	% of stomachs containing major (>4%) food items; N=772; yr-round; goldfish pond, Richland; see citation for detailed list of prey items and seasonal changes in diet
ar	% weight of an item; obtained by dividing the weight of a specific item by the total weight of all items; obtained from stomach contents; N=129; June - Nov.; ponds, Hanover County; % weight should be regarded with caution as it may represent a fraction of the weight of the specimen eaten. Mean weight of gut contents = 1.53 g
as	% occurrence of item; obtained by dividing # of a specific item taken by total # of items; based on stomach contents; N=129; June - Nov.; ponds, Hanover County; see citation for seasonal and body size variation in dietary composition
at	Percentage of bullfrog diet items; N=23-46; June - Aug.; Turkey Marsh, Kellogg Biological Station, Kalamazoo County; See paper for relationship between prey size and frog snout-vent length
au	% occurrence of 44 food items in stomach contents (species level); N=139; Mar. - June; streams, ponds, strip pits in Pulaski, Loanoke and Saline counties
av	Percent occurrence in stomach contents from frogs collected during day and night; N=124; ponds in north central TX
aw	% of stomachs in which food item was present at least once, frog weight range 10 - 200 g; N=300; San Joaquin Experimental Range, O'Neals; See paper for length to weight relationship
ax	Percent stomachs containing item; N=415; June - Nov.; ponds, Comanche and Kiowa counties
ay	% volume of stomach contents; N=18; summer; Breathitt and Perry counties
az	mean % body weight consumed per day, body weight mean = 55.8 g; N=25; Age=4 mo post-transformation; frogs fed a 1:1:1 mixture of mosquitofish, crickets and earthworms
ba	mean % body weight consumed per day, body weight mean = 29.8 g; N=25; Age=2 mo post-transformation; Frogs fed a 1:1:1 mixture of mosquitofish, crickets and earthworms
bb	mean % body weight consumed per day, body weight mean = 17.5 g; N=25; Age=1 mo post-transformation; Frogs fed a 1:1:1 mixture of mosquitofish, crickets and earthworms
bc	mean % body weight consumed per day, body weight mean = 42.4 g; N=25; Age=3 mo post-transformation; Frogs fed a 1:1:1 mixture of mosquitofish, crickets and earthworms
bd	average weight of stomach contents for frogs with an average body weight of 17.5 g; N=78; June - Nov.; ponds, Hanover County
be	average weight of stomach contents for frogs with an average body weight of 248.7 g; N=18; June - Nov.; ponds, Hanover County
bf	average weight of stomach contents for frogs with an average body weight of 106.3 g; N=42; June - Nov.; ponds, Hanover County
bg	N=34; Age=1 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bh	N=16; Age=0 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bi	N=44; Age=2 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bj	N=14; Age=3 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bk	N=10; Age=4 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bl	N=4; Age=6 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bm	N=4; Age=7 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]

bn	N=5; Age=5 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bo	N=46; Age=2 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bp	N=12; Age=3 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bq	N=7; Age=4 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
br	N=4; Age=5 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bs	N=4; Age=6 yr; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
bt	N=22; Crane Pond, E.S. George Reserve; See paper for growth rate vs. body length relationship
bu	see citation for review of growth rates based on size estimates; N=NR
bv	measured at 25C; Taylor-Kollros stages; N=11-14; Age=Taylor-Kollros stages 10-22; June - Aug.; see citation for temporal changes in body weights and effect of photoperiod on growth rate
bw	measured at 25C; Taylor-Kollros stages; N=11-14; Age=Taylor-Kollros stages 10-22; Nov. - Dec.; see citation for temporal changes in body weights and effect of photoperiod on growth rate
bx	mean activity radius; median = 7.2 ft; mode = 5.0-5.9 ft; N=131; Aug. - Sept.; pond at Go Home Bay, Ontario [lat., 45N; long., 79°o59'W]
by	distance of pond shoreline used; N=31; June - Sept.; farm ponds, Boone County; See citation for notes on occasional (<8% of frogs) pond to pond movement (distance traveled, 0.1 - 0.75 miles).
bz	figure of breathing frequency (breaths/min) at various seasons, temperatures and oxygen levels; N=6/group
ca	Maximum estimated age; N=567; Algonquin Provincial Park, Ontario [lat., 45°o35'N; long., 78°o30'W]
cb	N=NR
cc	N=NR; Edwin S. George Reserve, Ann Arbor
cd	figure of oxygen consumption at various body temperatures within seasons; N=6/group
ce	Oxygen consumption measured at 20C, frogs acclimated at 20C, mean body weight = 621 g (30.5 SE); N=8; See citation for relationship between temperature and oxygen consumption
cf	Oxygen consumption measured at 5C, frogs acclimated at 5C, mean body weight = 646 g (25.5 SE); N=8; See citation for relationship between temperature and oxygen consumption
cg	Mean total oxygen uptake at 20C (summed from skin and lung uptake), body weight = 228.2 g +/- 70.3 SE; N=6; See paper for carbon dioxide exchange rates
ch	Cutaneous oxygen uptake at 5C; N=9
ci	Pulmonary oxygen uptake at 5C; N=9
cj	Pulmonary oxygen uptake at 15C; N=4; See citation for relationship between oxygen uptake, body weight and temperature
ck	Cutaneous oxygen uptake at 15C; N=4; See citation for relationship between oxygen uptake, body weight and temperature
cl	Mean total oxygen uptake (skin + gills + lungs) measured at 25C, tadpole mean body weight = 5.74 g; N=5; See paper for relationship between body size, temperature and oxygen uptake
cm	Mean total oxygen uptake at 20C (summed from skin and gill uptake), aquatic tadpole (stage IV-V) body weight = 3.6 g +/- 1.03 SE; N=6; See paper for carbon dioxide exchange rates
cn	Mean total oxygen uptake at 20C (summed from skin and gill/lung uptake), air-breathing tadpole (stage XVI-XIX) body weight = 5.3 g +/- 1.2 SE; N=8; See paper for carbon dioxide exchange rates
co	figures of aerial activity oxygen consumption and body weights at different developmental stages; N=59
cq	figures of oxygen consumption of submerged tadpoles measured at various oxygen tensions and pre-metamorphic stages; N=10/group
cq	resting oxygen consumption at 23C; N=19; tadpoles (mean body weight = 1.96 g) collected from a reference site
cr	N=131; Aug. - Sept.; pond at Go Home Bay, Ontario [lat., 45N; long., 79°o59'W]
cs	mean density measured on both banks of river; N=3 visual counts; July - Sept.; lower Colorado River between Laguna and Morelos Dams; see citation for snout-vent length frequency diagrams (159 frogs, 45-180 mm) and influence of vegetation on population density
ct	N=20 quadrats; Age="overwintered" tadpoles; July; Angelo Coast Range Reserve [lat., 39°o44'N; long., 123°o39'W]
cu	N=17 quadrats; Age="overwintered" tadpoles; June; Angelo Coast Range Reserve [lat., 39°o44'N; long., 123°o39'W]
cv	N=3 sampling periods, 20,000 animals; Coldstream Pond and Fred Pond, northeast of Lexington, [38o06'23"N, 84o30'08"W]
cw	SA = Total Surface Area in cm <sup>2</sup> ; BW = body weight in grams wet weight; Log base 10; surface area estimated from removed skin of frogs ranging from 40 - 442 g; N=16
cx	Area of removed skin measured, equation based on weight (W) in grams; N=NR; See citation for graphical representation of data
cy	Mean total skin surface area measured by removing skin; air-breathing tadpole body weight 4.48 g +/- 0.31 SE; N=20
cz	probability of survival (age x-1 to age x); N=NR; Age=2 or 3 yrs; Edwin S. George Reserve
da	Non-breeding season survival; measured as # that returned to breed/# known to be alive at end of previous breeding season; N=25 - 54; Crane Pond, E.S. George Reserve
db	probability of survival (age x-1 to age x); N=NR; Age=5 yrs; Edwin S. George Reserve
dc	probability of survival (age x-1 to age x); N=NR; Age=4 yrs; Edwin S. George Reserve
dd	Range of mean survival of larvae; N=3 sampling periods, 20,000 animals; Nov. - July; May - Aug.; Sept. - June; Coldstream Pond and Fred Pond, northeast of Lexington, [38o06'23"N, 84o30'08"W]
de	Average distance between male frog and closest male neighbor within a chorus; also reported as a minimum territory radius of 9 ft.; N=94; June; Pond in E.S. George Reserve
df	Mean distance between territorial males; N=31; May; Basking Pond, Great Swamp National Wildlife Refuge, Morris County
dg	N=NR; Reviews time to transformation of bullfrog tadpoles
dh	N=NR; Ithaca
di	N=NR; Coldstream Pond and Fred Pond, northeast of Lexington, [38o06'23"N, 84o30'08"W]

dj	metamorphosis period for first breeding clutch; N=NR; lower Colorado River between Laguna and Morelos Dams
dk	time to metamorphosis for second clutch; N=NR; lower Colorado River between Laguna and Morelos Dams
dl	N=NR; Crane Pond, E.S. George Reserve, Livingston County [lat., 42.5N]; See paper for annual variation in percentage in metamorphic climax
dm	Tadpole transformation determined by presence of Taylor-Kollros stages XIX - XXIV; N=3500
dn	N=NR; farm ponds, Boone County
do	N=NR; Reviews breeding season of bullfrogs
dp	N=NR; Ridge Lake, Charleston, Coles County
dq	N=36; May; Basking Pond, Great Swamp National Wildlife Refuge, Morris County
dr	Measured as examining percentage of females with gonads in different developmental stages; N=350; farm ponds, Boone County
ds	laying; N=NR; Angelo Coast Range Reserve [lat., 39°o44'N; long., 123°o39'W]
dt	breeding season; N=100; lower Colorado River between Laguna and Morelos Dams
du	N=NR; farm ponds, Boone County
dv	N=6; Seiberling Naturealm, Summit County; see citation for overwintering activity monitoring via radiotelemetry
dw	N=NR; Ridge Lake, Charleston, Coles County
dx	"drinking" measured as 125I-iothalamate uptake in gut after 2 hrs; N=8

References

1 Cohen, Nathan W. and Walter E. Howard. 1958. Bullfrog food and growth at the San Joaquin Experimental Range, California. *Copeia*. 1958:223-225.

2 Ryan, Richard A. 1953. Growth rates of some ranids under natural conditions. *Copeia*. 1953(2):73-80.

3 Cecil, Stephen G. and John J. Just. 1979. Survival rate, population density and development of a naturally occurring anuran larvae (*Rana catesbeiana*). *Copeia*. 1979:447-453.

4 Clarkson, Robert W. and James C. deVos, Jr. 1986. The bullfrog, *Rana catesbeiana* Shaw, in the lower Colorado River, Arizona-California. *J. Herpetol.* 20:42-49.

5 Collins, James P. 1979. Intrapopulation variation in the body size at metamorphosis and timing of metamorphosis in the bullfrog. *Ecology*. 60(4):738-749.

6 Durham, Leonard and George W. Bennett. 1963. Age, growth, and homing in the bullfrog. *J. Wildl. Manage.* 27:107-123.

7 Viparina, Sally and John J. Just. 1975. The life period, growth and differentiation of *Rana catesbeiana* larvae occurring in nature. *Copeia*. 1975:103-109.

8 Howard, Richard D. 1981. Sexual dimorphism in bullfrogs. *Ecology*. 62(2):303-310.

9 Shirose, L.J., R.J. Brooks, J.R. Barta and S.S. Desser. 1993. Intersexual differences in growth, mortality, and size at maturity in bullfrogs in central Ontario. *Can. J. Zool.* 71(12):2363-2369.

10 Rocha, Priscila L. and Luiz G.S. Branco. 1998. Seasonal changes in the cardiovascular, respiratory and metabolic responses to temperature and hypoxia in the bullfrog *Rana catesbeiana*. *J. Exp. Biol.* 201:761-768.

11 Weathers, Wesley W. 1976. Influence of temperature acclimation on oxygen consumption, haemodynamics and oxygen transport in bullfrogs. *Aust. J. Zool.* 24:321-330.

12 McAlpine, Donald F. and Timothy G. Dilworth. 1989. Microhabitat and prey size among three species of *Rana* (Anura: Ranidae) sympatric in eastern Canada. *Can. J. Zool.* 67:2244-2252.

13 Modzelewski, Edward H., Jr. and Dudley D. Culley, Jr. 1974. Growth responses of the bullfrog, *Rana catesbeiana*, fed various live foods. *Herpetologica*. 30(4):396-405.

14 McKamie, J.A. and G.A. Heidt. 1974. A comparison of spring food habits of the bullfrog, *Rana catesbeiana*, in three habitats of central Arkansas. *Southwest. Nat.* 19(1):107-111.

15 Ryan, Michael J. 1980. The reproductive behavior of the bullfrog (*Rana catesbeiana*). *Copeia*. 1980:108-114.

16 Howard, Richard D. 1978. The influence of male-defended oviposition sites on early embryo mortality in bullfrogs. *Ecology*. 59(4):789-798.

17 Emlen, Stephen T. 1977. "Double clutching" and its possible significance in the bullfrog. *Copeia*. 1977(4):749-751.

18 Bury, R. B. and J.A. Whelan. 1984. Ecology and management of the bullfrog. Washington D.C.: U.S. Department of Interior, Fish and Wildlife Service, Resource Publication No. 155. 23 p.

19 Stuart, James N. and Charles W. Painter. 1993. *Rana catesbeiana* (bullfrog) cannibalism. *Herpetologica*. 24(3):103.

20 Stewart, Margaret M. and Peggy Sandison. 1972. Comparative food habits of sympatric mink frogs, bullfrogs, and green frogs. *J. Herpetol.* 6(3/4):241-244.

21 Corse, Wayne A. and Dean E. Metter. 1980. Economics, adult feeding and larval growth of *Rana catesbeiana* on a fish hatchery. *J. Herpetol.* 14:231-238.

22 Brooks, G. R., Jr. 1964. An analysis of the food habits of the bullfrog by body size, sex, month, and habitat. *Va. J. Sci.* 15:173-186.

23 Werner, Earl E., Gary A. Wellborn and Mark A. McPeck. 1995. Diet composition in postmetamorphic bullfrogs and green frogs: implications for interspecific predation and competition. *J. Herpetol.* 29(4):600-607.

24 Carpenter, H. L. and Morrison, E. O. 1973. Feeding behavior of the bullfrog, *Rana catesbeiana*, in north central Texas. *Bios.* 44:188-193.

25 Tyler, Jack. D. and Roger D. Hoestenbach, Jr. 1979. Differences in food of bullfrogs (*Rana catesbeiana*) from pond and stream habitats in southwestern Oklahoma. *Southwest. Nat.* 24(1):33-38.

26 Bush, Francis M. 1959. Foods of some Kentucky herptiles. *Herpetologica*. 15:73-77.

27 Turner, F.B. 1960. Postmetamorphic growth in anurans. *Am. Midl. Nat.* 64:327-338.

28 Crawshaw, Larry I.; Richard N. Rausch; Lonnie P. Wollmuth and Ethan J. Bauer. 1992. Seasonal rhythms of development and temperature selection in larval bullfrogs, *Rana catesbeiana* Shaw. *Physiol. Zool.* 65(2):346-359.

29 Currie, William and Edward D. Bellis. 1969. Home range and movements of the bullfrog, *Rana catesbeiana* (Shaw), in an Ontario pond. *Copeia*. 1969:688-692.

30 Willis, Yuell L., Don L. Moyle, and Thomas S. Baskett. 1956. Emergence, breeding, hibernation, movements and transformation of the bullfrog, *Rana catesbeiana*, in Missouri. *Copeia*. 1956:30-41.

31 Storer, Tracy I. 1922. The eastern bullfrog in California. *Calif. Fish Game.* 8:219-224.



- 32 Howard, Richard D. 1978. The evolution of mating strategies in bullfrogs, *Rana catesbeiana*. *Evolution*. 32(4):850-874.
- 33 Burggren, Warren W., and Nigel H. West. 1982. Changing respiratory importance of gills, lungs and skin during metamorphosis in the bullfrog *Rana catesbeiana*. *Respir. Physiol.* 47:151-164.
- 34 Hutchinson, Victor H., W.G. Whitford and M. Kohl. 1968. Relation of body size and surface area to gas exchange in anurans. *Physiol. Zool.* 41:65-85.
- 35 Burggren, Warren W., Martin E. Feder, and Alan W. Pinder. 1983. Temperature and the balance between aerial and aquatic respiration in larvae of *Rana berlandieri* and *Rana catesbeiana*. *Physiol. Zool.* 56(2):263-273.
- 36 Hillman, Stanley S. and Malcom S. Lea. 1983. Aerial activity oxygen consumption during metamorphosis of the bullfrog, *Rana catesbeiana*. *Copeia*. 1983(2):407-410.
- 37 Crowder, William C., Minghua Nie and Gordon R. Ultsch. 1998. Oxygen uptake in bullfrog tadpoles (*Rana catesbeiana*). *J. Exp. Zool.* 280:121-134.
- 38 Rowe, Christopher L., Owen M. Kinney, Roy D. Nagle and Justin D. Congdon. 1998. Elevated maintenance costs in an anuran (*Rana catesbeiana*) exposed to a mixture of trace elements during the embryonic and early larval periods. *Physiol. Zool.* 71:27-35.
- 39 Kupferberg, Sarah J. 1997. Bullfrog (*Rana catesbeiana*) invasion of a California river: the role of larval competition. *Ecology*. 78:1736-1751.
- 40 Talbot, Colleen R. and Martin E. Feder. 1992. Relationships among cutaneous surface area, cutaneous mass, and body mass in frogs: A reappraisal. *Physiol. Zool.* 65(6):1135-1147.
- 41 Howard, Richard D. 1984. Alternative mating behaviors of young male bullfrogs. *Am. Zool.* 24:397-406.
- 42 Emlen, Stephen T. 1968. Territoriality in the bullfrog. *Copeia*. 1968(2):240-243.
- 43 Stinner, John, Nicholas Zarlinga and Scott Orcutt. 1994. Overwintering behavior of adult bullfrogs, *Rana catesbeiana*, in northeastern Ohio. *Ohio J. Sci.* 94(1):8-13.
- 44 Bentley, P.J. and T. Yorio. 1979. Do frogs drink?. *J. Exp. Biol.* 79:41-46.

\*Cal/EPA, OEHHA and the University of California Regents are not responsible for damages of any kind resulting from the use of or reliance on information in this report. Users are encouraged to consult the original data. Updated: February 1999.